



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Hypoglycaemia associated with gastrointestinal and extragastrointestinal stromal tumour in two dogs

Citation for published version:

Lodzinska, J, Ballber, C, Smith, S & Bavcar, S 2019, 'Hypoglycaemia associated with gastrointestinal and extragastrointestinal stromal tumour in two dogs', *Vet Record Case Reports*, vol. 7, no. 4.
<https://doi.org/10.1136/vetreccr-2018-000784>

Digital Object Identifier (DOI):

[10.1136/vetreccr-2018-000784](https://doi.org/10.1136/vetreccr-2018-000784)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Vet Record Case Reports

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



VetRecordCaseReports

HYPOGLYCAEMIA ASSOCIATED WITH GASTROINTESTINAL AND EXTRAGASTROINTESTINAL STROMAL TUMOUR IN TWO DOGS.

Journal:	<i>Veterinary Record Case Reports</i>
Manuscript ID	vetreccr-2018-000784.R2
Manuscript Type:	Companion or pet animals
Species:	Dogs
Date Submitted by the Author:	23-Oct-2019
Complete List of Authors:	Lodzinska, Joanna; University of Edinburgh Royal Dick School of Veterinary Studies, Ballber, Clara; Woodstreet Veterinary Hospital Smith, Sionagh; University of Edinburgh, Veterinary Pathology Unit Bavcar, Spela; University of Edinburgh Royal Dick School of Veterinary Studies
Keywords:	Dogs, Computed tomography (CT), Ultrasonography, GIST, EGIST
Topics:	Radiology, Oncology, Emergency medicine and critical care
Abstract:	Gastrointestinal stromal tumours (GISTs) are uncommon mesenchymal tumours that originate from the interstitial cells of Cajal (ICC). As these tumours are difficult to distinguish from gastrointestinal smooth muscle tumours using standard histological techniques, their true prevalence may be underestimated. Metabolic and systemic consequences of GISTs are not well described in any species. More rarely, neoplasms with histological and immunohistochemical features similar to GISTs may occur outside the gastrointestinal tract, so-called Extra-gastrointestinal Stromal Tumours (EGIST). EGISTs have never been described in the veterinary literature. In this article we present and describe clinical findings, management and treatment of two dogs with clinical hypoglycaemia induced by histologically confirmed GIST and EGIST. Hypoglycaemia resolved immediately and long-term after tumour excision. To the authors' knowledge this is the first report of hypoglycaemia associated with a canine GIST and the first case report of an EGIST in the dog.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Veterinary Record casereports

Submission template for full cases

All case reports MUST be submitted online using this Word template. Please refer to the [Hints and Tips template](#) so you know what to include in each section

- o <http://mc.manuscriptcentral.com/vetreccr>
- o You will be asked for more detailed information on submission where you can also upload images, multimedia files, etc
- o Further details are available in the [Instructions for authors](#)

For studies using client-owned animals the article must demonstrate a high standard (best practice) of veterinary care and have informed client consent

You, your practice or institution must be a subscriber to Veterinary Record Case Reports in order to submit. Subscribers can submit as many cases as they like, access all the published material, and re-use any published material for personal use and teaching without further permission.

- o For more information on rates and how to purchase your subscription visit <http://vetrecordcasereports.bmj.com/site/about/Subslanding.xhtml>
- o Contact your librarian, head of department or practice owner to see if your institution already has a subscription

TITLE OF CASE <i>Do not include "a case report"</i>
HYPOGLYCAEMIA ASSOCIATED WITH GASTROINTESTINAL AND EXTRAGASTROINTESTINAL STROMAL TUMOUR IN TWO DOGS.
SUMMARY <i>Up to 150 words summarising the case presentation and outcome (this will be freely available online)</i>
Gastrointestinal stromal tumours (GISTs) are uncommon mesenchymal tumours that originate from the interstitial cells of Cajal (ICC). As these tumours are difficult to distinguish from gastrointestinal smooth muscle tumours using standard histological techniques, their true prevalence may be underestimated. Metabolic and systemic consequences of GISTs are not well described in any species. More rarely, neoplasms with histological and immunohistochemical features similar to GISTs may occur outside the gastrointestinal tract, so-called Extra-gastrointestinal Stromal Tumours (EGIST). EGISTs have never been described in the veterinary literature. In this article we present and describe clinical findings, management and treatment of two dogs with clinical hypoglycaemia induced by histologically confirmed GIST and EGIST.

Hypoglycaemia resolved immediately and long-term after tumour excision. To the authors' knowledge this is the first report of hypoglycaemia associated with a canine GIST and the first case report of an EGIST in the dog.

BACKGROUND *Why you think this case is important – why did you write it up?*

GISTs are mesenchymal tumours that occur in the GIT of dogs (1, 2), cats (3, 4), humans (5-8), horses (9) and non-human primates (10). They can occur in any segment of the GIT, however most of them develop in the stomach and small intestine. Clinically, they can lead to focal obstruction, vomiting and diarrhoea as well as chronic low grade GIT blood loss, resulting in iron deficiency anaemia. Surgical removal usually results in excellent outcomes (11) while only a few case reports have been published demonstrating a response to tyrosine kinase inhibitors (12-14).

More rarely, neoplasms with histological and immunohistochemical features similar to GISTs may occur outside the gastrointestinal tract, so-called Extra-gastrointestinal Stromal Tumours (EGISTs).

These tumours are difficult to distinguish from gastrointestinal smooth muscle tumours such as leiomyoma or leiomyosarcoma (11) using standard histological techniques, therefore their true prevalence may be underestimated. It has been established that GISTs can be differentiated by the expression of KIT (CD117), a receptor tyrosine kinase encoded by the proto-oncogene *c-KIT* (11, 15). One study also suggests that the immunohistochemical application of DOG1 (discovered on GIST protein 1) achieves higher specificity and sensitivity than KIT for differentiating between canine GISTs and leiomyosarcomas. Inclusion of both DOG1 and KIT IHC in diagnostic panels offers improved overall accuracy of canine GIST diagnosis (16).

No previous cases of GIST-associated hypoglycaemia have been described in the veterinary literature. To our knowledge there are also no case reports describing an EGIST in veterinary medicine.

Hypoglycaemia in dogs, defined as a blood glucose concentration of less than 3.3mmol/L (17), can be a life-threatening disorder that necessitates swift and decisive treatment to prevent lasting

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

neuronal damage. Most frequently reported causes of severe hypoglycaemia in adult dogs are insulinoma, excessive insulin administration, and non-pancreatic neoplasia (18, 19). Sepsis, severe hepatopathy, and hypoadrenocorticism are usually associated with only a mild decrease in blood glucose concentration (18).

In human medicine, non-islet cell tumour hypoglycaemia (NICTH) is a well-recognised paraneoplastic syndrome (20) and has been described in association with gastrointestinal stromal tumours (GISTs) in humans (21). In dogs, NICTH has been described with neoplasms of hepatic (22, 23), smooth muscle (24, 25), mammary (26) and renal origin (27, 28), but not with neoplasms of the gastrointestinal tract (GIT). Different mechanisms have been proposed to explain the paraneoplastic syndrome. NICTH may be caused by increased tumour utilization of glucose; decreased hepatic glycogenolysis or gluconeogenesis; or secretion of insulin or insulin-like growth factor 1 (IGF1) and insulin-like growth factor 2 (IGF2)(29) . However, in human medicine it has been established that IGF2 is the main cause of NICTH, including in GISTs (30). Two different studies suggest that IGF2 is also a relevant factor in dogs (25, 26). Some mesenchymal and epithelial tumours can overexpress IGF2, resulting in the secretion of partially processed precursors of IGF2 ('big' IGF2). This molecule binds to insulin receptors, resulting in insulin-like hypoglycaemic effects (31). The purpose of this paper is to describe clinical findings and treatment in two dogs with marked hypoglycaemia associated with GIST and EGIST that resolved after tumour excision.

CASE PRESENTATION *Presenting features, clinical and environmental history*

DOG 1

A 13-year-old, male intact Golden retriever canine presented to the referring emergency service with a history of intermittent diarrhoea and acute onset weakness and collapse. Symptomatic hypoglycaemia (2.4-2.6 mmol/l; reference range 4.11-7.95 mmol/l) causing weakness episodes was detected by the referring veterinary surgeon on multiple measurements and the dog was referred to the Royal (Dick) School of Veterinary Studies for investigation of hypoglycaemic

episodes and a palpable cranial abdominal mass. The initial treatment by the referring veterinarian included 2.5% glucose constant rate infusion. On physical examination the dog was quiet, alert and responsive. Ptyalism was noted and a large mass was palpated in the cranial abdomen. Initial investigations included complete blood count and serum biochemistry that revealed low urea and confirmed hypoglycaemia of 2.58 mmol/l (reference range 4.11-7.95 mmol/l). Despite glucose supplementation, the glucose was consistently low at 2.4 mmol/l and 2.6 mmol/l on subsequent measurements taken over a 24-hour period using a hand-held glucometer. To rule out hepatic dysfunction as cause of low urea and glucose, a bile acid stimulation test was performed, and this ruled out severe functional loss.

Abdominal ultrasound scan confirmed the presence of a large round heterogeneous mass containing small cavities in the right cranial abdomen dorsal to the duodenum (Fig. 1). Its exact origin could not be established but neoplasia was considered the most likely differential diagnosis. No other abdominal abnormalities were identified (including in the liver) and thoracic radiography did not show any evidence of metastatic disease or any other abnormality. The patient was stabilised on glucose infusion and surgery was scheduled.

Ventral midline exploratory coeliotomy was performed. The mass was associated with the junction of the central and right pancreatic regions and pancreatic origin was suspected but not surgically confirmed, given multiple adhesions to the duodenum, gastric wall and transverse colon. The common bile duct and the gastric pylorus were not involved. No other abnormalities were found during surgery and, given the normal liver morphology and lack of significant laboratory changes, liver biopsies were not taken. The mass was successfully excised and submitted for histopathology. Glucose supplementation was gradually discontinued 12 hours after the surgery. The hypoglycaemia resolved quickly and all the measurements during the post-operative period were within normal limits. The dog was managed on pain relief and discharged four days later.

Macroscopically, the mass was fluctuant to firm, mottled dark purple to grey and spherical, measuring 14cm diameter, with fat attached to its surface. On cut section, it was solid and mottled

pink, red and orange, with a soft consistency. Histopathology of the mass confirmed an intra-abdominal malignant spindle cell tumour (sarcoma) that was completely excised with narrow margins and no evidence of pancreatic or GIT involvement. The only recognisable architecture consisted of fat and smooth muscle. This was effaced by a well-demarcated, expansile and encapsulated proliferation of spindle cells arranged in interlacing fascicles with a faint herringbone pattern. The spindle cells breached the capsule and invaded the surrounding fat. The spindle cells contained fibrillar, eosinophilic cytoplasm and elongated or “cigar shaped” nuclei. Anisokaryosis was mild and the mitotic index averaged 3 per hpf at 400X. There were multifocal areas of necrosis and haemorrhage, with occasional cystic degeneration. Immunohistochemically, ~10% of the spindle cells labelled strongly with antibody against CD117, confirming a diagnosis of EGIST.

In addition to surgical removal, treatment in this patient consisted of single-agent chemotherapy (four doses of doxorubicin (30mg/m²) given every 3 weeks). Six months after the initial diagnosis a new 8mm nodule next to the pancreas was noticed on a routine recheck ultrasound. Fine needle aspirate (FNA) confirmed mesenchymal neoplasia, indicating probable recurrence of the previously diagnosed EGIST. Blood glucose levels were tested and were within normal limits on subsequent chemotherapy consultations. Five months later multiple pancreatic nodules and a liver mass were identified on ultrasound (Fig. 2). The owners opted for no further treatment and the dog was humanely euthanised at the referring practice.

DOG 2

A 13-year-old, female neutered Labrador canine was referred for investigation of an abdominal mass. A few episodes of vomiting and diarrhoea were reported but they had resolved by the time of presentation. The owner also reported a 2-month history of progressive weight loss. On physical examination the patient was bright alert and responsive with a body condition score of 2/9 (32). A grade I/VI heart murmur was auscultated and abdominal palpation indicated the presence of a mass in the cranial abdomen. Initial investigations included complete blood count and serum biochemistry. Mild leukopenia with mild neutropenia (no left shift) and lymphopenia,

1 and minimal (non-significant) normocytic-normochromic anaemia were noted. Biochemical
2 abnormalities consisted of moderate elevations in both alanine transaminase (ALT) and alkaline
3 phosphatase (AP). Mildly raised basal bile acids were not considered significant, but a bile acid
4 stimulation test was not performed. Due to the non-specific presenting complaints and mild
5 hypercholesterolemia, TT4 and TSH were assessed but were not suggestive of hypothyroidism.
6

7
8 Abdominal ultrasound confirmed the presence of a very large cranial abdominal mass, for which
9 the exact origin could not be established. Results of an FNA suggested chronic haemorrhage.
10
11 Computed tomography (CT) of the abdomen indicated that the mass was most likely of gastric
12 origin. Additional findings on thoracic CT images showed changes consistent with aspiration
13 pneumonia, though there was no history or clinical signs typical of that. Elective surgery was
14 scheduled and the patient was discharged on the same day with a course of antibiotics to treat the
15 presumed subclinical aspiration pneumonia prior to surgery. The dog re-presented to the
16 R(D)SVS emergency service a few hours later due to a collapsing episode. The dog was dull but
17 responsive, with continued ptyalism. In addition, pyrexia and tachypnoea with increased
18 inspiratory effort had developed. The heart rate was 104 beats per minute, with good quality
19 synchronous pulses, and blood pressure was 155mmHg. Blood analysis revealed more severe
20 lymphopenia and neutropenia with a glucose of 4.3 mmol/l (reference range 4.11-7.95 mmol/l).
21
22 Oxygen therapy and intravenous antibiotic therapy were initiated and the patient was closely
23 monitored. Two hours later the glucose levels had dropped to 3mmol/l. At this point a dextrose
24 bolus was administered and repeat (every 2 hours) glucose measurements using hand-held
25 glucometer were commenced. These were consistently low therefore a 5% dextrose continuous
26 rate infusion (CRI) was started.
27

28
29 The respiratory status of the patient remained stable in the following days but glucose remained
30 low despite CRI supplementation. Administration of an anti-inflammatory dose of dexamethasone
31 achieved a mild improvement in glucose levels. However, they decreased again once
32 dexamethasone was discontinued. Leukocytes returned to normal, although lymphopenia
33 persisted. Once the respiratory status of the patient was stable, a ventral midline exploratory
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

coeliotomy was performed. It revealed a mass arising from the dorsal aspect of the pyloric antrum with omentum wrapped around it; this was removed and submitted for histopathology. Glucose levels were closely monitored during surgery and due to initial hypoglycaemia of 2.8mmol/l, supplemental dextrose injections were given. Once the mass was excised the consecutive glucose readings were within normal limits and no further supplementation was required. Recovery from the surgery was uneventful. The hypoglycaemia resolved completely after the mass excision and the patient was discharged four days later.

Macroscopically, the mass was ovoid and multinodular, measuring 14cm diameter at its widest dimension, with an attached fragment of omentum. Histopathologically it comprised an expansile, non-encapsulated but well demarcated proliferation of spindle cells similar to those described above (Fig.3). Anisokaryosis was moderate to marked but mitoses were rare. There were large areas of necrosis and some haemorrhage. The spindle cell proliferation merged with the muscularis layer of the pyloric stomach. Immunohistochemically, 20% of the spindle cells labelled moderately strongly with antibody against CD117, supporting a diagnosis of GIST (Fig.4). Different chemotherapy protocols were discussed but declined by the owners. At the last follow up appointment (three months after initial diagnosis) the dog was free of clinical signs.

INVESTIGATIONS *If relevant*

DIFFERENTIAL DIAGNOSIS *If relevant*

1	TREATMENT <i>If relevant</i>
2	
3	
4	
5	
6	
7	OUTCOME AND FOLLOW-UP
8	
9	
10	
11	
12	DISCUSSION <i>Include a very brief review of similar published cases</i>
13	
14	As far as the authors are aware, there are no previous reports of GIST-associated hypoglycaemia
15	in the dog. To date, only one case report of GIST-associated hypoglycaemia in a veterinary
16	patient has been published, and that was in a horse (33). However, GISTs are a relatively new
17	entity. An immunohistochemical review of previously diagnosed gastrointestinal leiomyomas and
18	leiomyosarcomas has resulted in most of these tumours being re-classified as GISTs (11, 15, 34).
19	All the cases describing hypoglycaemia secondary to leiomyomas/sarcomas (19, 24, 25, 35, 36)
20	were published prior to this new classification. Therefore, it is possible that some of these
21	previous cases were actually GISTs.
22	
23	Paraneoplastic syndromes are common manifestations of many malignancies and have been
24	reported with GIST (37, 38). Multiple articles on GIST-associated hypoglycaemia are available in
25	human medicine (39-42) but no information on its systemic effects is available in the veterinary
26	literature. Hypoglycaemia was a common presenting abnormality in both of the dogs reported
27	here. In dog 1, the hypoglycaemia was considered to be the most likely cause of the clinical signs
28	(weakness and collapse), given the lack of other physical and laboratory abnormalities. However,
29	other unrelated causes (such as a transient arrhythmia or syncopal episode) were not ruled out.
30	Liver dysfunction was considered an unlikely explanation for the hypoglycaemia. A bile acid
31	stimulation test was not consistent with severe liver dysfunction and no imaging abnormalities
32	were identified in the liver. Furthermore, the dog became normoglycaemic after the tumour was
33	resected, providing more support for a paraneoplastic mechanism.
34	
35	During surgery the mass was adjacent to the pancreas, but pancreatic involvement was not
36	confirmed histopathologically. A concomitant, undetected insulinoma cannot be ruled out as a

potential cause for hypoglycaemia as insulin levels were not measured but the presence of two different neoplasms is considered unlikely.

Interestingly, histopathological examination of the mass in dog 1 did not indicate a direct connection between the mass and the GIT. EGISTs have been described in human medicine widely but nothing has been reported in the veterinary literature. In recent years, ICC or ICC-like cells have been found in extraintestinal organs (43). For example, in the genitourinary tract, KIT-positive specialized pacemaker cells have been described in the human bladder, urethra, uterus, and prostate (44-48). The presence of these cells in various organs or soft tissue could provide an explanation for the rare cases of GISTs that occur as primary tumours outside the GIT, such as in the mesentery, omentum, retroperitoneum, liver, gallbladder, vagina, uterus, urinary bladder, or prostate (44, 49-58). In some of these tumours, the origin remains unclear (59).

In addition to its hypoglycaemia, dog 2 was also diagnosed with aspiration pneumonia and, considering the clinical findings (dyspnoea, pyrexia), laboratory abnormalities (neutropenia), and timeframe, sepsis with secondary hypoglycaemia was a possibility. However, hypoglycaemia persisted after sepsis resolved. It also responded to glucocorticoid administration and, as with case 1, resolved after mass removal, such that sepsis as a sole cause was unlikely.

The main limitation of this report is the fact that the exact underlying mechanism for the hypoglycaemia was not determined. Given that the main cause of NICTH seems to be the secretion of IGF2, this should ideally have been measured to confirm the source of hypoglycaemia. Consequently, other causes of hypoglycaemia (insulinoma or liver disease in case 1 and sepsis in case 2), although unlikely, cannot be completely excluded. Furthermore, while an antibody panel was not applied in these two cases, both samples labelled strongly with antibody against CD117 which, given the other histological features, was considered sufficiently diagnostic of GIST and EGIST.

To the authors' knowledge these are the first cases describing hypoglycaemia associated with GIST and EGIST in the dog. GIST and EGIST should be considered as a differential diagnosis in dogs presenting with an abdominal mass and hypoglycaemia.

LEARNING POINTS/TAKE HOME MESSAGES 3 to 5 bullet points – this is a required field

- EGISTs do occur in dogs.
- GIST and EGIST should be considered as a differential diagnosis in dogs presenting with an abdominal mass.
- GIST and EGIST should be considered as a differential diagnosis in dogs presenting hypoglycaemia.

REFERENCES *Vancouver style*

1. Gillespie V, Baer K, Farrelly J, Craft D, Luong R. Canine gastrointestinal stromal tumors: immunohistochemical expression of CD34 and examination of prognostic indicators including proliferation markers Ki67 and AgNOR. *Veterinary pathology*. 2011;48(1):283-91.
2. LaRock RG, Ginn PE. Immunohistochemical staining characteristics of canine gastrointestinal stromal tumors. *Veterinary pathology*. 1997;34(4):303-11.
3. Morini M, Gentilini F, Pietra M, Spadari A, Turba ME, Mandrioli L, et al. Cytological, immunohistochemical and mutational analysis of a gastric gastrointestinal stromal tumour in a cat. *Journal of comparative pathology*. 2011;145(2-3):152-7.
4. Suwa A, Shimoda T. Intestinal gastrointestinal stromal tumor in a cat. *J Vet Med Sci*. 2017;79(3):562-6.
5. Gupta A, Gupta S, Tandon A, Kotru M, Kumar S. Gastrointestinal stromal tumor causing ileo-ileal intussusception in an adult patient a rare presentation with review of literature. *The Pan African Medical Journal*. 2011;8:29.
6. Kim CJ, Day S, Yeh KA. Gastrointestinal stromal tumors: analysis of clinical and pathologic factors. *The American surgeon*. 2001;67(2):135-7.
7. Occhionorelli S, Mitaritunno M, Pennella A, Carella M, Presicce L, Pellegrini D, et al. Gastro-intestinal stromal tumor (GIST): case report. *Il Giornale di chirurgia*. 2001;22(3):65-9.
8. Zhao X, Yue C. Gastrointestinal stromal tumor. *Journal of Gastrointestinal Oncology*. 2012;3(3):189-208.

9. Wong D, Hepworth K, Yaeger M, Miles K, Wilgenbusch C. Imaging diagnosis-hypoglycemia associated with cholangiocarcinoma and peritoneal carcinomatosis in a horse. *Vet Radiol Ultrasound*. 2015;56(1):E9-12.
10. Saturday GA, Lasota J, Frost D, Brasky KB, Hubbard G, Miettinen M. KIT-positive gastrointestinal stromal tumor in a 22-year-old male chimpanzee (*Pan troglodites*). *Veterinary pathology*. 2005;42(3):362-5.
11. Russell KN, Mehler SJ, Skorupski KA, Baez JL, Shofer FS, Goldschmidt MH. Clinical and immunohistochemical differentiation of gastrointestinal stromal tumors from leiomyosarcomas in dogs: 42 cases (1990-2003). *J Am Vet Med Assoc*. 2007;230(9):1329-33.
12. Elliott JW, Swinbourne F, Parry A, Baines L. Successful treatment of a metastatic, gastrointestinal stromal tumour in a dog with toceranib phosphate (Palladia). *J Small Anim Pract*. 2017;58(7):416-8.
13. Irie M, Takeuchi Y, Ohtake Y, Suzuki H, Nagata N, Miyoshi T, et al. Imatinib mesylate treatment in a dog with gastrointestinal stromal tumors with a c-kit mutation. *J Vet Med Sci*. 2015;77(11):1535-9.
14. Kobayashi M, Kuroki S, Ito K, Yasuda A, Sawada H, Ono K, et al. Imatinib-associated tumour response in a dog with a non-resectable gastrointestinal stromal tumour harbouring a c-kit exon 11 deletion mutation. *Veterinary journal (London, England : 1997)*. 2013;198(1):271-4.
15. Maas CPHJ, Ter Haar G, Van Der Gaag I, Kirpensteijn J. Reclassification of Small Intestinal and Cecal Smooth Muscle Tumors in 72 Dogs: Clinical, Histologic, and Immunohistochemical Evaluation. *Veterinary Surgery*. 2007;36(4):302-13.
16. Dailey DD, Ehrhart EJ, Duval DL, Bass T, Powers BE. DOG1 is a sensitive and specific immunohistochemical marker for diagnosis of canine gastrointestinal stromal tumors. *Journal of veterinary diagnostic investigation : official publication of the American Association of Veterinary Laboratory Diagnosticians, Inc*. 2015;27(3):268-77.
17. Ettinger SJ, Feldman EC, Côté E. *Textbook of veterinary internal medicine : diseases of the dog and the cat*. Eighth edition.. ed: St. Louis, Missouri : Elsevier; 2017.

18. Leifer CE, Peterson ME. Hypoglycemia. 1984. p. 873-89.
19. Leifer CE, Peterson ME, Matus RE, Patnaik AK. Hypoglycemia associated with nonislet cell tumor in 13 dogs. *J Am Vet Med Assoc*. 1985;186(1):53-5.
20. Bodnar TW, Acevedo MJ, Pietropaolo M. Management of Non-Islet-Cell Tumor Hypoglycemia: A Clinical Review. *The Journal of Clinical Endocrinology & Metabolism*. 2014;99(3):713-22.
21. Beckers MMJ, Slee PHTJ, Van Doorn J. Hypoglycaemia in a patient with a gastrointestinal stromal tumour. *Clinical Endocrinology*. 2003;59(3):402-4.
22. Battaglia L, Petterino C, Zappulli V, Castagnaro M. Hypoglycaemia as a paraneoplastic syndrome associated with renal adenocarcinoma in a dog. *Veterinary research communications*. 2005;29(8):671-5.
23. Zini E, Glaus TM, Minuto F, Arvigo M, Hauser B, Reusch CE. Paraneoplastic hypoglycemia due to an insulin-like growth factor type-II secreting hepatocellular carcinoma in a dog. *J Vet Intern Med*. 2007;21(1):193-5.
24. Beaudry D, Knapp DW, Montgomery T, Sandusky GS, Morrison WB, Nelson RW. Hypoglycemia in four dogs with smooth muscle tumors. *J Vet Intern Med*. 1995;9(6):415-8.
25. Boari A, Barreca A, Bestetti GE, Minuto F, Venturoli M. Hypoglycemia in a dog with a leiomyoma of the gastric wall producing an insulin-like growth factor II-like peptide. *European journal of endocrinology*. 1995;132(6):744-50.
26. Rossi G, Errico G, Perez P, Rossi G, Paltrinieri S. Paraneoplastic hypoglycemia in a diabetic dog with an insulin growth factor-2-producing mammary carcinoma. *Vet Clin Pathol*. 2010;39(4):480-4.
27. Swain JM, Pirie RS, Hudson NP, Else RW, Evans H, McGorum BC. Insulin-like growth factors and recurrent hypoglycemia associated with renal cell carcinoma in a horse. *J Vet Intern Med*. 2005;19(4):613-6.
28. Snead EC. A case of bilateral renal lymphosarcoma with secondary polycythaemia and paraneoplastic syndromes of hypoglycaemia and uveitis in an English Springer Spaniel.

Veterinary and Comparative Oncology. 2005;3(3):139-44.

29. Withrow SJ, MacEwan EG. Small animal clinical oncology. Third edition.. ed. Philadelphia: Philadelphia : W. B. Saunders; 2001.

30. Rikhof B, Van Den Berg G, Van Der Graaf WTA. Non-islet cell tumour hypoglycaemia in a patient with a gastrointestinal stromal tumour. Acta Oncologica, 2005, Vol44(7), p764-766. 2005;44(7):764-6.

31. de Groot JW, Rikhof B, van Doorn J, Bilo HJ, Alleman MA, Honkoop AH, et al. Non-islet cell tumour-induced hypoglycaemia: a review of the literature including two new cases. Endocrine-related cancer. 2007;14(4):979-93.

32. WSAVA nutritional assessment guidelines. Journal of feline medicine and surgery. 2011;13(7):516-25.

33. Haga HA, Ytrehus B, Rudshaug IJ, Ottesen N. Gastrointestinal stromal tumour and hypoglycemia in a Fjord pony: case report. Acta veterinaria Scandinavica. 2008;50:9.

34. Frost D, Lasota J, Miettinen M. Gastrointestinal stromal tumors and leiomyomas in the dog: a histopathologic, immunohistochemical, and molecular genetic study of 50 cases. Veterinary pathology. 2003;40(1):42-54.

35. Bagley RS, Levy JK, Malarkey DE. Hypoglycemia associated with intra-abdominal leiomyoma and leiomyosarcoma in six dogs. J Am Vet Med Assoc. 1996;208(1):69-71.

36. Cohen M, Post GS, Wright JC. Gastrointestinal leiomyosarcoma in 14 dogs. J Vet Intern Med. 2003;17(1):107-10.

37. Guirgis HM, Holcombe RF. A case of advanced gastrointestinal stromal tumor (GIST) presenting with paraneoplastic syndrome. Journal of Clinical Oncology. 2004;22(14_suppl):4242-.

38. Tsikrikas S, Manolakopoulos S, Deutsch M, Alexakis G, Sialevis K, Giannopoulos D, et al. Unusual combination of paraneoplastic manifestations in a patient with metastatic gastrointestinal stromal tumor (GIST). Scandinavian journal of gastroenterology. 2008;43(8):1012-5.

39. Dimitriadis GK, Gopalakrishnan K, Rao R, Grammatopoulos DK, Randeva HS, Weickert MO, et al. Severe paraneoplastic hypoglycemia secondary to a gastrointestinal stromal tumour masquerading as a stroke. *Endocrinology, diabetes & metabolism case reports*. 2015;2015:150062.
40. Escobar GA, Robinson WA, Nydam TL, Heiple DC, Weiss GJ, Buckley L, et al. Severe paraneoplastic hypoglycemia in a patient with a gastrointestinal stromal tumor with an exon 9 mutation: a case report. *BMC cancer*. 2007;7:13.
41. Guiteau J, Fanucchi M, Folpe A, Staley CA, 3rd, Kooby DA. Hypoglycemia in the setting of advanced gastrointestinal stromal tumor. *The American surgeon*. 2006;72(12):1225-30.
42. Singhal A, Hadi R, Mehrotra K, Rastogi S, Masood S. Paraneoplastic Hypoglycaemia: A Rare Manifestation of Pelvic Gastrointestinal Stromal Tumour. *Journal of clinical and diagnostic research : JCDR*. 2017;11(2):Xd01-xd2.
43. Huizinga JD, Faussone-Pellegrini MS. About the presence of interstitial cells of Cajal outside the musculature of the gastrointestinal tract. *Journal of cellular and molecular medicine*. 2005;9(2):468-73.
44. Ciontea SM, Radu E, Regalia T, Ceafalan L, Cretoiu D, Gherghiceanu M, et al. C-kit immunopositive interstitial cells (Cajal-type) in human myometrium. *Journal of cellular and molecular medicine*. 2005;9(2):407-20.
45. Piaseczna Piotrowska A, Rolle U, Solari V, Puri P. Interstitial cells of Cajal in the human normal urinary bladder and in the bladder of patients with megacystis-microcolon intestinal hypoperistalsis syndrome. *BJU international*. 2004;94(1):143-6.
46. Shafik A, El-Sibai O, Shafik AA, Shafik I. Identification of interstitial cells of Cajal in human urinary bladder: concept of vesical pacemaker. *Urology*. 2004;64(4):809-13.
47. Van der Aa F, Roskams T, Blyweert W, De Ridder D. Interstitial cells in the human prostate: a new therapeutic target? *The Prostate*. 2003;56(4):250-5.
48. van der AF, Roskams T, Blyweert W, Ost D, Bogaert G, De Ridder D. Identification of kit positive cells in the human urinary tract. *The Journal of urology*. 2004;171(6 Pt 1):2492-6.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
49. Dickson BC, Srigley JR, Pollett AF, Blackstein ME, Honey JD, Juco JW. Rectal gastrointestinal stromal tumor mimicking a primary prostatic lesion. *The Canadian journal of urology*. 2008;15(3):4112-4.
50. Furihata M, Fujimori T, Imura J, Ono Y, Furihata T, Shimoda M, et al. Malignant stromal tumor, so called "gastrointestinal stromal tumor", with rhabdomyomatous differentiation occurring in the gallbladder. *Pathology, research and practice*. 2005;201(8-9):609-13.
51. Herawi M, Montgomery EA, Epstein JI. Gastrointestinal stromal tumors (GISTs) on prostate needle biopsy: A clinicopathologic study of 8 cases. *The American journal of surgical pathology*. 2006;30(11):1389-95.
52. Krska Z, Peskova M, Povysil C, Horejs J, Sedlackova E, Kudrnova Z. GIST of pancreas. *Prague medical report*. 2005;106(2):201-8.
53. Lee CH, Lin YH, Lin HY, Lee CM, Chu JS. Gastrointestinal stromal tumor of the prostate: a case report and literature review. *Human pathology*. 2006;37(10):1361-5.
54. Madden JF, Burchette JL, Raj GV, Daly JT, Tannenbaum M. Anterior rectal wall gastrointestinal stromal tumor presenting clinically as prostatic mass. *Urologic oncology*. 2005;23(4):268-72.
55. Reith JD, Goldblum JR, Lyles RH, Weiss SW. Extragastrointestinal (soft tissue) stromal tumors: an analysis of 48 cases with emphasis on histologic predictors of outcome. *Modern pathology : an official journal of the United States and Canadian Academy of Pathology, Inc*. 2000;13(5):577-85.
56. Van der Aa F, Sciort R, Blyweert W, Ost D, Van Poppel H, Van Oosterom A, et al. Gastrointestinal stromal tumor of the prostate. *Urology*. 2005;65(2):388.
57. Voelzke BB, Sakamoto K, Hantel A, Paner GP, Kash J, Waters WB, et al. Gastrointestinal stromal tumor: involvement in urologic patients and recent therapeutic advances. *Urology*. 2002;60(2):218-22.
58. Wingen CB, Pauwels PA, Debiec-Rychter M, van Gemert WG, Vos MC. Uterine gastrointestinal stromal tumour (GIST). *Gynecologic oncology*. 2005;97(3):970-2.

59. Takahashi RH, Matsubayashi J, Yokotsuka M, Tachibana M, Kusama H, Nagao T. An intrapelvic extraintestinal gastrointestinal stromal tumor of undetermined origin: diagnosis by prostate needle biopsy. Pathology, research and practice. 2012;208(12):736-40.

FIGURE/VIDEO CAPTIONS *figures should NOT be embedded in this document*

Fig. 1

Ultrasonographic image from dog 1 showing large round heterogeneous mass (between callipers) with multiple hypoechoic cavitations (asterisk shows the largest one) which was identified in the right cranial abdomen dorsal to the duodenum.

Fig. 2

Ultrasonographic image from dog 1 showing multiple different sized, hypoechoic nodules within the pancreatic parenchyma (between callipers) discovered five months after initial presentation. Note hepatic parenchyma in the near field (margin delineated with arrows).

Fig. 3

Abdominal gastrointestinal stromal tumour from dog 2 [HE]. The mass comprises a partially necrotic proliferation of moderately pleomorphic spindle cells arranged in interlacing streams.

Fig. 4

Immunohistochemistry of abdominal gastrointestinal stromal tumour from dog 2. Spindle cells label moderately strongly with antibody against CD117.

OWNER'S PERSPECTIVE *Optional*

Copyright Statement

I, Joanna Lodzinska, The Corresponding Author, has the right to assign on behalf of all authors and does assign on behalf of all authors, a full assignment of all intellectual property rights for all content within the submitted case report (other than as agreed with the BMJ Publishing Group Ltd and the British Veterinary Association) ("BMJ" and "BVA") in any media known now or created in the future, and permits this case report (if accepted) to be published on Veterinary Record Case Reports and to be fully exploited within the remit of the assignment as set out in the assignment which has been read <http://journals.bmj.com/site/misc/vetreccrcopyright.pdf>

Date: 29/11/2018

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

<p>PLEASE SAVE YOUR TEMPLATE WITH THE FOLLOWING FORMAT:</p> <p>Corresponding author’s last name and date of submission, eg,</p> <p>Smith_June_2017.doc</p>

Veterinary Record casereports

Submission template for full cases

All case reports MUST be submitted online using this Word template. Please refer to the [Hints and Tips template](#) so you know what to include in each section

- <http://mc.manuscriptcentral.com/vetreccr>
- You will be asked for more detailed information on submission where you can also upload images, multimedia files, etc
- Further details are available in the [Instructions for authors](#)

For studies using client-owned animals the article must demonstrate a high standard (best practice) of veterinary care and have informed client consent

You, your practice or institution must be a subscriber to Veterinary Record Case Reports in order to submit. Subscribers can submit as many cases as they like, access all the published material, and re-use any published material for personal use and teaching without further permission.

- For more information on rates and how to purchase your subscription visit <http://vetrecordcasereports.bmj.com/site/about/Subslanding.xhtml>
- Contact your librarian, head of department or practice owner to see if your institution already has a subscription

TITLE OF CASE *Do not include "a case report"*

HYPOGLYCAEMIA ASSOCIATED WITH GASTROINTESTINAL AND EXTRAGASTROINTESTINAL STROMAL TUMOUR IN TWO DOGS.

SUMMARY *Up to 150 words summarising the case presentation and outcome (this will be freely available online)*

Gastrointestinal stromal tumours (GISTs) are uncommon mesenchymal tumours that originate from the interstitial cells of Cajal (ICC). As these tumours are difficult to distinguish from gastrointestinal smooth muscle tumours using standard histological techniques, their true prevalence may be underestimated. Metabolic and systemic consequences of GISTs are not well described in any species. More rarely, neoplasms with histological and immunohistochemical features similar to GISTs may occur outside the gastrointestinal tract, so-called Extra-gastrointestinal Stromal Tumours (EGIST). EGISTs have never been described in the veterinary literature. In this article we present and describe clinical findings, management and treatment of two dogs with clinical hypoglycaemia induced by histologically confirmed GIST and EGIST.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Hypoglycaemia resolved immediately and long-term after tumour excision. To the authors' knowledge this is the first report of hypoglycaemia associated with a canine GIST and the first case report of an EGIST in the dog.

BACKGROUND *Why you think this case is important – why did you write it up?*

GISTs are mesenchymal tumours that occur in the GIT of dogs (1, 2), cats (3, 4), humans (5-8), horses (9) and non-human primates (10). They can occur in any segment of the GIT, however most of them develop in the stomach and small intestine. Clinically, they can lead to focal obstruction, vomiting and diarrhoea as well as chronic low grade GIT blood loss, resulting in iron deficiency anaemia. Surgical removal usually results in excellent outcomes (11) while only a few case reports have been published demonstrating a response to tyrosine kinase inhibitors (12-14). More rarely, neoplasms with histological and immunohistochemical features similar to GISTs may occur outside the gastrointestinal tract, so-called Extra-gastrointestinal Stromal Tumours (EGISTs).

These tumours are difficult to distinguish from gastrointestinal smooth muscle tumours such as leiomyoma or leiomyosarcoma (11) using standard histological techniques, therefore their true prevalence may be underestimated. It has been established that GISTs can be differentiated by the expression of KIT (CD117), a receptor tyrosine kinase encoded by the proto-oncogene *c-KIT* (11, 15). One study also suggests that the immunohistochemical application of DOG1 (discovered on GIST protein 1) achieves higher specificity and sensitivity than KIT for differentiating between canine GISTs and leiomyosarcomas. Inclusion of both DOG1 and KIT IHC in diagnostic panels offers improved overall accuracy of canine GIST diagnosis (16).

No previous cases of GIST-associated hypoglycaemia have been described in the veterinary literature. To our knowledge there are also no case reports describing an EGIST in veterinary medicine.

Hypoglycaemia in dogs, defined as a blood glucose concentration of less than 3.3mmol/L (17), can be a life-threatening disorder that necessitates swift and decisive treatment to prevent lasting

neuronal damage. Most frequently reported causes of severe hypoglycaemia in adult dogs are insulinoma, excessive insulin administration, and non-pancreatic neoplasia (18, 19). Sepsis, severe hepatopathy, and hypoadrenocorticism are usually associated with only a mild decrease in blood glucose concentration (18).

In human medicine, non-islet cell tumour hypoglycaemia (NICTH) is a well-recognised paraneoplastic syndrome (20) and has been described in association with gastrointestinal stromal tumours (GISTs) in humans (21). In dogs, NICTH has been described with neoplasms of hepatic (22, 23), smooth muscle (24, 25), mammary (26) and renal origin (27, 28), but not with neoplasms of the gastrointestinal tract (GIT). Different mechanisms have been proposed to explain the paraneoplastic syndrome. NICTH may be caused by increased tumour utilization of glucose; decreased hepatic glycogenolysis or gluconeogenesis; or secretion of insulin or insulin-like growth factor 1 (IGF1) and insulin-like growth factor 2 (IGF2)(29). However, in human medicine it has been established that IGF2 is the main cause of NICTH, including in GISTs (30). Two different studies suggest that IGF2 is also a relevant factor in dogs (25, 26). Some mesenchymal and epithelial tumours can overexpress IGF2, resulting in the secretion of partially processed precursors of IGF2 ('big' IGF2). This molecule binds to insulin receptors, resulting in insulin-like hypoglycaemic effects (31). The purpose of this paper is to describe clinical findings and treatment in two dogs with marked hypoglycaemia associated with GIST and EGIST that resolved after tumour excision.

CASE PRESENTATION *Presenting features, clinical and environmental history*

DOG 1

A 13-year-old, male intact Golden retriever canine presented to the referring emergency service with a history of intermittent diarrhoea and acute onset weakness and collapse. Symptomatic hypoglycaemia (2.4-2.6 mmol/l; reference range 4.11-7.95 mmol/l) causing weakness episodes was detected by the referring veterinary surgeon on multiple measurements and the dog was referred to the Royal (Dick) School of Veterinary Studies for investigation of hypoglycaemic

episodes and a palpable cranial abdominal mass. The initial treatment by the referring veterinarian included 2.5% glucose constant rate infusion. On physical examination the dog was quiet, alert and responsive. Ptyalism was noted and a large mass was palpated in the cranial abdomen. Initial investigations included complete blood count and serum biochemistry that revealed low urea and confirmed hypoglycaemia of 2.58 mmol/l (reference range 4.11-7.95 mmol/l). Despite glucose supplementation, the glucose was consistently low at 2.4 mmol/l and 2.6 mmol/l on subsequent measurements taken over a 24-hour period using a hand-held glucometer. To rule out hepatic dysfunction as cause of low urea and glucose, a bile acid stimulation test was performed, and this ruled out severe functional loss.

Abdominal ultrasound scan confirmed the presence of a large round heterogeneous mass containing small cavities in the right cranial abdomen dorsal to the duodenum (Fig. 1). Its exact origin could not be established but neoplasia was considered the most likely differential diagnosis. No other abdominal abnormalities were identified (including in the liver) and thoracic radiography did not show any evidence of metastatic disease or any other abnormality. The patient was stabilised on glucose infusion and surgery was scheduled.

Ventral midline exploratory coeliotomy was performed. The mass was associated with the junction of the central and right pancreatic regions and pancreatic origin was suspected but not surgically confirmed, given multiple adhesions to the duodenum, gastric wall and transverse colon. The common bile duct and the gastric pylorus were not involved. No other abnormalities were found during surgery and, given the normal liver morphology and lack of significant laboratory changes, liver biopsies were not taken. The mass was successfully excised and submitted for histopathology. Glucose supplementation was gradually discontinued 12 hours after the surgery. The hypoglycaemia resolved quickly and all the measurements during the post-operative period were within normal limits. The dog was managed on pain relief and discharged four days later.

Macroscopically, the mass was fluctuant to firm, mottled dark purple to grey and spherical, measuring 14cm diameter, with fat attached to its surface. On cut section, it was solid and mottled

pink, red and orange, with a soft consistency. Histopathology of the mass confirmed an intra-abdominal malignant spindle cell tumour (sarcoma) that was completely excised with narrow margins and no evidence of pancreatic or GIT involvement. The only recognisable architecture consisted of fat and smooth muscle. This was effaced by a well-demarcated, expansile and encapsulated proliferation of spindle cells arranged in interlacing fascicles with a faint herringbone pattern. The spindle cells breached the capsule and invaded the surrounding fat. The spindle cells contained fibrillar, eosinophilic cytoplasm and elongated or “cigar shaped” nuclei. Anisokaryosis was mild and the mitotic index averaged 3 per hpf at 400X. There were multifocal areas of necrosis and haemorrhage, with occasional cystic degeneration. Immunohistochemically, ~10% of the spindle cells labelled strongly with antibody against CD117, confirming a diagnosis of EGIST.

In addition to surgical removal, treatment in this patient consisted of single-agent chemotherapy (four doses of doxorubicin (30mg/m²) given every 3 weeks). Six months after the initial diagnosis a new 8mm nodule next to the pancreas was noticed on a routine recheck ultrasound. Fine needle aspirate (FNA) confirmed mesenchymal neoplasia, indicating probable recurrence of the previously diagnosed EGIST. Blood glucose levels were tested and were within normal limits on subsequent chemotherapy consultations. Five months later multiple pancreatic nodules and a liver mass were identified on ultrasound (Fig. 2). The owners opted for no further treatment and the dog was humanely euthanised at the referring practice.

DOG 2

A 13-year-old, female neutered Labrador canine was referred for investigation of an abdominal mass. A few episodes of vomiting and diarrhoea were reported but they had resolved by the time of presentation. The owner also reported a 2-month history of progressive weight loss. On physical examination the patient was bright alert and responsive with a body condition score of 2/9 (32). A grade I/VI heart murmur was auscultated and abdominal palpation indicated the presence of a mass in the cranial abdomen. Initial investigations included complete blood count and serum biochemistry. Mild leukopenia with mild neutropenia (no left shift) and lymphopenia,

and minimal (non-significant) normocytic-normochromic anaemia were noted. Biochemical abnormalities consisted of moderate elevations in both alanine transaminase (ALT) and alkaline phosphatase (AP). Mildly raised basal bile acids were not considered significant, but a bile acid stimulation test was not performed. Due to the non-specific presenting complaints and mild hypercholesterolemia, TT4 and TSH were assessed but were not suggestive of hypothyroidism.

Abdominal ultrasound confirmed the presence of a very large cranial abdominal mass, for which the exact origin could not be established. Results of an FNA suggested chronic haemorrhage. Computed tomography (CT) of the abdomen indicated that the mass was most likely of gastric origin. Additional findings on thoracic CT images showed changes consistent with aspiration pneumonia, though there was no history or clinical signs typical of that. Elective surgery was scheduled and the patient was discharged on the same day with a course of antibiotics to treat the presumed subclinical aspiration pneumonia prior to surgery. The dog re-presented to the R(D)SVS emergency service a few hours later due to a collapsing episode. The dog was dull but responsive, with continued ptyalism. In addition, pyrexia and tachypnoea with increased inspiratory effort had developed. The heart rate was 104 beats per minute, with good quality synchronous pulses, and blood pressure was 155mmHg. Blood analysis revealed more severe lymphopenia and neutropenia with a glucose of 4.3 mmol/l (reference range 4.11-7.95 mmol/l). Oxygen therapy and intravenous antibiotic therapy were initiated and the patient was closely monitored. Two hours later the glucose levels had dropped to 3mmol/l. At this point a dextrose bolus was administered and repeat (every 2 hours) glucose measurements using hand-held glucometer were commenced. These were consistently low therefore a 5% dextrose continuous rate infusion (CRI) was started.

The respiratory status of the patient remained stable in the following days but glucose remained low despite CRI supplementation. Administration of an anti-inflammatory dose of dexamethasone achieved a mild improvement in glucose levels. However, they decreased again once dexamethasone was discontinued. Leukocytes returned to normal, although lymphopenia persisted. Once the respiratory status of the patient was stable, a ventral midline exploratory

coeliotomy was performed. It revealed a mass arising from the dorsal aspect of the pyloric antrum with omentum wrapped around it; this was removed and submitted for histopathology. Glucose levels were closely monitored during surgery and due to initial hypoglycaemia of 2.8mmol/l, supplemental dextrose injections were given. Once the mass was excised the consecutive glucose readings were within normal limits and no further supplementation was required. Recovery from the surgery was uneventful. The hypoglycaemia resolved completely after the mass excision and the patient was discharged four days later.

Macroscopically, the mass was ovoid and multinodular, measuring 14cm diameter at its widest dimension, with an attached fragment of omentum. Histopathologically it comprised an expansile, non-encapsulated but well demarcated proliferation of spindle cells similar to those described above (Fig.3). Anisokaryosis was moderate to marked but mitoses were rare. There were large areas of necrosis and some haemorrhage. The spindle cell proliferation merged with the muscularis layer of the pyloric stomach. Immunohistochemically, 20% of the spindle cells labelled moderately strongly with antibody against CD117, supporting a diagnosis of GIST (Fig.4). Different chemotherapy protocols were discussed but declined by the owners. At the last follow up appointment (three months after initial diagnosis) the dog was free of clinical signs.

INVESTIGATIONS *If relevant***DIFFERENTIAL DIAGNOSIS *If relevant***

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TREATMENT <i>If relevant</i>
OUTCOME AND FOLLOW-UP
DISCUSSION <i>Include a very brief review of similar published cases</i>
<p>As far as the authors are aware, there are no previous reports of GIST-associated hypoglycaemia in the dog. To date, only one case report of GIST-associated hypoglycaemia in a veterinary patient has been published, and that was in a horse (33). However, GISTs are a relatively new entity. An immunohistochemical review of previously diagnosed gastrointestinal leiomyomas and leiomyosarcomas has resulted in most of these tumours being re-classified as GISTs (11, 15, 34). All the cases describing hypoglycaemia secondary to leiomyomas/sarcomas (19, 24, 25, 35, 36) were published prior to this new classification. Therefore, it is possible that some of these previous cases were actually GISTs.</p> <p>Paraneoplastic syndromes are common manifestations of many malignancies and have been reported with GIST (37, 38). Multiple articles on GIST-associated hypoglycaemia are available in human medicine (39-42) but no information on its systemic effects is available in the veterinary literature. Hypoglycaemia was a common presenting abnormality in both of the dogs reported here. In dog 1, the hypoglycaemia was considered to be the most likely cause of the clinical sings (weakness and collapse), given the lack of other physical and laboratory abnormalities. However, other unrelated causes (such as a transient arrhythmia or syncopal episode) were not ruled out. Liver dysfunction was considered an unlikely explanation for the hypoglycaemia. A bile acid stimulation test was not consistent with severe liver dysfunction and no imaging abnormalities were identified in the liver. Furthermore, the dog became normoglycaemic after the tumour was resected, providing more support for a paraneoplastic mechanism.</p> <p>During surgery the mass was adjacent to the pancreas, but pancreatic involvement was not confirmed histopathologically. A concomitant, undetected insulinoma cannot be ruled out as a</p>

1 potential cause for hypoglycaemia as insulin levels were not measured but the presence of two
2 different neoplasms is considered unlikely.
3

4 Interestingly, histopathological examination of the mass in dog 1 did not indicate a direct
5 connection between the mass and the GIT. EGISTs have been described in human medicine
6 widely but nothing has been reported in the veterinary literature. In recent years, ICC or ICC-like
7 cells have been found in extraintestinal organs (43). For example, in the genitourinary tract, KIT-
8 positive specialized pacemaker cells have been described in the human bladder, urethra, uterus,
9 and prostate (44-48). The presence of these cells in various organs or soft tissue could provide an
10 explanation for the rare cases of GISTs that occur as primary tumours outside the GIT, such as in
11 the mesentery, omentum, retroperitoneum, liver, gallbladder, vagina, uterus, urinary bladder, or
12 prostate (44, 49-58). In some of these tumours, the origin remains unclear (59).
13
14

15 In addition to its hypoglycaemia, dog 2 was also diagnosed with aspiration pneumonia and,
16 considering the clinical findings (dyspnoea, pyrexia), laboratory abnormalities (neutropenia), and
17 timeframe, sepsis with secondary hypoglycaemia was a possibility. However, hypoglycaemia
18 persisted after sepsis resolved. It also responded to glucocorticoid administration and, as with
19 case 1, resolved after mass removal, such that sepsis as a sole cause was unlikely.
20
21

22 The main limitation of this report is the fact that the exact underlying mechanism for the
23 hypoglycaemia was not determined. Given that the main cause of NICTH seems to be the
24 secretion of IGF2, this should ideally have been measured to confirm the source of
25 hypoglycaemia. Consequently, other causes of hypoglycaemia (insulinoma or liver disease in
26 case 1 and sepsis in case 2), although unlikely, cannot be completely excluded. Furthermore,
27 while an antibody panel was not applied in these two cases, both samples labelled strongly with
28 antibody against CD117 which, given the other histological features, was considered sufficiently
29 diagnostic of GIST and EGIST.
30
31

32 To the authors' knowledge these are the first cases describing hypoglycaemia associated with
33 GIST and EGIST in the dog. GIST and EGIST should be considered as a differential diagnosis in
34 dogs presenting with an abdominal mass and hypoglycaemia.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

LEARNING POINTS/TAKE HOME MESSAGES 3 to 5 bullet points – this is a required field

- EGISTs do occur in dogs.
- GIST and EGIST should be considered as a differential diagnosis in dogs presenting with an abdominal mass.
- GIST and EGIST should be considered as a differential diagnosis in dogs presenting hypoglycaemia.

REFERENCES *Vancouver style*

1. Gillespie V, Baer K, Farrelly J, Craft D, Luong R. Canine gastrointestinal stromal tumors: immunohistochemical expression of CD34 and examination of prognostic indicators including proliferation markers Ki67 and AgNOR. *Veterinary pathology*. 2011;48(1):283-91.
2. LaRock RG, Ginn PE. Immunohistochemical staining characteristics of canine gastrointestinal stromal tumors. *Veterinary pathology*. 1997;34(4):303-11.
3. Morini M, Gentilini F, Pietra M, Spadari A, Turba ME, Mandrioli L, et al. Cytological, immunohistochemical and mutational analysis of a gastric gastrointestinal stromal tumour in a cat. *Journal of comparative pathology*. 2011;145(2-3):152-7.
4. Suwa A, Shimoda T. Intestinal gastrointestinal stromal tumor in a cat. *J Vet Med Sci*. 2017;79(3):562-6.
5. Gupta A, Gupta S, Tandon A, Kotru M, Kumar S. Gastrointestinal stromal tumor causing ileo-ileal intussusception in an adult patient a rare presentation with review of literature. *The Pan African Medical Journal*. 2011;8:29.
6. Kim CJ, Day S, Yeh KA. Gastrointestinal stromal tumors: analysis of clinical and pathologic factors. *The American surgeon*. 2001;67(2):135-7.
7. Occhionorelli S, Mitaritunno M, Pennella A, Carella M, Presicce L, Pellegrini D, et al. Gastro-intestinal stromal tumor (GIST): case report. *Il Giornale di chirurgia*. 2001;22(3):65-9.
8. Zhao X, Yue C. Gastrointestinal stromal tumor. *Journal of Gastrointestinal Oncology*. 2012;3(3):189-208.

9. Wong D, Hepworth K, Yaeger M, Miles K, Wilgenbusch C. Imaging diagnosis-hypoglycemia associated with cholangiocarcinoma and peritoneal carcinomatosis in a horse. *Vet Radiol Ultrasound*. 2015;56(1):E9-12.
10. Saturday GA, Lasota J, Frost D, Brasky KB, Hubbard G, Miettinen M. KIT-positive gastrointestinal stromal tumor in a 22-year-old male chimpanzee (*Pan troglodites*). *Veterinary pathology*. 2005;42(3):362-5.
11. Russell KN, Mehler SJ, Skorupski KA, Baez JL, Shofer FS, Goldschmidt MH. Clinical and immunohistochemical differentiation of gastrointestinal stromal tumors from leiomyosarcomas in dogs: 42 cases (1990-2003). *J Am Vet Med Assoc*. 2007;230(9):1329-33.
12. Elliott JW, Swinbourne F, Parry A, Baines L. Successful treatment of a metastatic, gastrointestinal stromal tumour in a dog with toceranib phosphate (Palladia). *J Small Anim Pract*. 2017;58(7):416-8.
13. Irie M, Takeuchi Y, Ohtake Y, Suzuki H, Nagata N, Miyoshi T, et al. Imatinib mesylate treatment in a dog with gastrointestinal stromal tumors with a c-kit mutation. *J Vet Med Sci*. 2015;77(11):1535-9.
14. Kobayashi M, Kuroki S, Ito K, Yasuda A, Sawada H, Ono K, et al. Imatinib-associated tumour response in a dog with a non-resectable gastrointestinal stromal tumour harbouring a c-kit exon 11 deletion mutation. *Veterinary journal (London, England : 1997)*. 2013;198(1):271-4.
15. Maas CPHJ, Ter Haar G, Van Der Gaag I, Kirpensteijn J. Reclassification of Small Intestinal and Cecal Smooth Muscle Tumors in 72 Dogs: Clinical, Histologic, and Immunohistochemical Evaluation. *Veterinary Surgery*. 2007;36(4):302-13.
16. Dailey DD, Ehrhart EJ, Duval DL, Bass T, Powers BE. DOG1 is a sensitive and specific immunohistochemical marker for diagnosis of canine gastrointestinal stromal tumors. *Journal of veterinary diagnostic investigation : official publication of the American Association of Veterinary Laboratory Diagnosticians, Inc*. 2015;27(3):268-77.
17. Ettinger SJ, Feldman EC, Côté E. *Textbook of veterinary internal medicine : diseases of the dog and the cat*. Eighth edition.. ed: St. Louis, Missouri : Elsevier; 2017.

18. Leifer CE, Peterson ME. Hypoglycemia. 1984. p. 873-89.
19. Leifer CE, Peterson ME, Matus RE, Patnaik AK. Hypoglycemia associated with nonislet cell tumor in 13 dogs. *J Am Vet Med Assoc.* 1985;186(1):53-5.
20. Bodnar TW, Acevedo MJ, Pietropaolo M. Management of Non-Islet-Cell Tumor Hypoglycemia: A Clinical Review. *The Journal of Clinical Endocrinology & Metabolism.* 2014;99(3):713-22.
21. Beckers MMJ, Slee PHTJ, Van Doorn J. Hypoglycaemia in a patient with a gastrointestinal stromal tumour. *Clinical Endocrinology.* 2003;59(3):402-4.
22. Battaglia L, Petterino C, Zappulli V, Castagnaro M. Hypoglycaemia as a paraneoplastic syndrome associated with renal adenocarcinoma in a dog. *Veterinary research communications.* 2005;29(8):671-5.
23. Zini E, Glaus TM, Minuto F, Arvigo M, Hauser B, Reusch CE. Paraneoplastic hypoglycemia due to an insulin-like growth factor type-II secreting hepatocellular carcinoma in a dog. *J Vet Intern Med.* 2007;21(1):193-5.
24. Beaudry D, Knapp DW, Montgomery T, Sandusky GS, Morrison WB, Nelson RW. Hypoglycemia in four dogs with smooth muscle tumors. *J Vet Intern Med.* 1995;9(6):415-8.
25. Boari A, Barreca A, Bestetti GE, Minuto F, Venturoli M. Hypoglycemia in a dog with a leiomyoma of the gastric wall producing an insulin-like growth factor II-like peptide. *European journal of endocrinology.* 1995;132(6):744-50.
26. Rossi G, Errico G, Perez P, Rossi G, Paltrinieri S. Paraneoplastic hypoglycemia in a diabetic dog with an insulin growth factor-2-producing mammary carcinoma. *Vet Clin Pathol.* 2010;39(4):480-4.
27. Swain JM, Pirie RS, Hudson NP, Else RW, Evans H, McGorum BC. Insulin-like growth factors and recurrent hypoglycemia associated with renal cell carcinoma in a horse. *J Vet Intern Med.* 2005;19(4):613-6.
28. Snead EC. A case of bilateral renal lymphosarcoma with secondary polycythaemia and paraneoplastic syndromes of hypoglycaemia and uveitis in an English Springer Spaniel.

Veterinary and Comparative Oncology. 2005;3(3):139-44.

29. Withrow SJ, MacEwan EG. Small animal clinical oncology. Third edition.. ed. Philadelphia: Philadelphia : W. B. Saunders; 2001.

30. Rikhof B, Van Den Berg G, Van Der Graaf WTA. Non-islet cell tumour hypoglycaemia in a patient with a gastrointestinal stromal tumour. Acta Oncologica, 2005, Vol44(7), p764-766. 2005;44(7):764-6.

31. de Groot JW, Rikhof B, van Doorn J, Bilo HJ, Alleman MA, Honkoop AH, et al. Non-islet cell tumour-induced hypoglycaemia: a review of the literature including two new cases. Endocrine-related cancer. 2007;14(4):979-93.

32. WSAVA nutritional assessment guidelines. Journal of feline medicine and surgery. 2011;13(7):516-25.

33. Haga HA, Ytrehus B, Rudshaug IJ, Ottesen N. Gastrointestinal stromal tumour and hypoglycemia in a Fjord pony: case report. Acta veterinaria Scandinavica. 2008;50:9.

34. Frost D, Lasota J, Miettinen M. Gastrointestinal stromal tumors and leiomyomas in the dog: a histopathologic, immunohistochemical, and molecular genetic study of 50 cases. Veterinary pathology. 2003;40(1):42-54.

35. Bagley RS, Levy JK, Malarkey DE. Hypoglycemia associated with intra-abdominal leiomyoma and leiomyosarcoma in six dogs. J Am Vet Med Assoc. 1996;208(1):69-71.

36. Cohen M, Post GS, Wright JC. Gastrointestinal leiomyosarcoma in 14 dogs. J Vet Intern Med. 2003;17(1):107-10.

37. Guirgis HM, Holcombe RF. A case of advanced gastrointestinal stromal tumor (GIST) presenting with paraneoplastic syndrome. Journal of Clinical Oncology. 2004;22(14_suppl):4242-.

38. Tsikrikas S, Manolakopoulos S, Deutsch M, Alexakis G, Sialevis K, Giannopoulos D, et al. Unusual combination of paraneoplastic manifestations in a patient with metastatic gastrointestinal stromal tumor (GIST). Scandinavian journal of gastroenterology. 2008;43(8):1012-5.

39. Dimitriadis GK, Gopalakrishnan K, Rao R, Grammatopoulos DK, Randeva HS, Weickert MO, et al. Severe paraneoplastic hypoglycemia secondary to a gastrointestinal stromal tumour masquerading as a stroke. *Endocrinology, diabetes & metabolism case reports*. 2015;2015:150062.
40. Escobar GA, Robinson WA, Nydam TL, Heiple DC, Weiss GJ, Buckley L, et al. Severe paraneoplastic hypoglycemia in a patient with a gastrointestinal stromal tumor with an exon 9 mutation: a case report. *BMC cancer*. 2007;7:13.
41. Guiteau J, Fanucchi M, Folpe A, Staley CA, 3rd, Kooby DA. Hypoglycemia in the setting of advanced gastrointestinal stromal tumor. *The American surgeon*. 2006;72(12):1225-30.
42. Singhal A, Hadi R, Mehrotra K, Rastogi S, Masood S. Paraneoplastic Hypoglycaemia: A Rare Manifestation of Pelvic Gastrointestinal Stromal Tumour. *Journal of clinical and diagnostic research : JCDR*. 2017;11(2):Xd01-xd2.
43. Huizinga JD, Faussone-Pellegrini MS. About the presence of interstitial cells of Cajal outside the musculature of the gastrointestinal tract. *Journal of cellular and molecular medicine*. 2005;9(2):468-73.
44. Ciontea SM, Radu E, Regalia T, Ceafalan L, Cretoiu D, Gherghiceanu M, et al. C-kit immunopositive interstitial cells (Cajal-type) in human myometrium. *Journal of cellular and molecular medicine*. 2005;9(2):407-20.
45. Piaseczna Piotrowska A, Rolle U, Solari V, Puri P. Interstitial cells of Cajal in the human normal urinary bladder and in the bladder of patients with megacystis-microcolon intestinal hypoperistalsis syndrome. *BJU international*. 2004;94(1):143-6.
46. Shafik A, El-Sibai O, Shafik AA, Shafik I. Identification of interstitial cells of Cajal in human urinary bladder: concept of vesical pacemaker. *Urology*. 2004;64(4):809-13.
47. Van der Aa F, Roskams T, Blyweert W, De Ridder D. Interstitial cells in the human prostate: a new therapeutic target? *The Prostate*. 2003;56(4):250-5.
48. van der AF, Roskams T, Blyweert W, Ost D, Bogaert G, De Ridder D. Identification of kit positive cells in the human urinary tract. *The Journal of urology*. 2004;171(6 Pt 1):2492-6.

49. Dickson BC, Srigley JR, Pollett AF, Blackstein ME, Honey JD, Juco JW. Rectal gastrointestinal stromal tumor mimicking a primary prostatic lesion. The Canadian journal of urology. 2008;15(3):4112-4.
50. Furihata M, Fujimori T, Imura J, Ono Y, Furihata T, Shimoda M, et al. Malignant stromal tumor, so called "gastrointestinal stromal tumor", with rhabdomyomatous differentiation occurring in the gallbladder. Pathology, research and practice. 2005;201(8-9):609-13.
51. Herawi M, Montgomery EA, Epstein JI. Gastrointestinal stromal tumors (GISTs) on prostate needle biopsy: A clinicopathologic study of 8 cases. The American journal of surgical pathology. 2006;30(11):1389-95.
52. Krska Z, Peskova M, Povysil C, Horejs J, Sedlackova E, Kudrnova Z. GIST of pancreas. Prague medical report. 2005;106(2):201-8.
53. Lee CH, Lin YH, Lin HY, Lee CM, Chu JS. Gastrointestinal stromal tumor of the prostate: a case report and literature review. Human pathology. 2006;37(10):1361-5.
54. Madden JF, Burchette JL, Raj GV, Daly JT, Tannenbaum M. Anterior rectal wall gastrointestinal stromal tumor presenting clinically as prostatic mass. Urologic oncology. 2005;23(4):268-72.
55. Reith JD, Goldblum JR, Lyles RH, Weiss SW. Extragastrointestinal (soft tissue) stromal tumors: an analysis of 48 cases with emphasis on histologic predictors of outcome. Modern pathology : an official journal of the United States and Canadian Academy of Pathology, Inc. 2000;13(5):577-85.
56. Van der Aa F, Sciort R, Blyweert W, Ost D, Van Poppel H, Van Oosterom A, et al. Gastrointestinal stromal tumor of the prostate. Urology. 2005;65(2):388.
57. Voelzke BB, Sakamoto K, Hantel A, Paner GP, Kash J, Waters WB, et al. Gastrointestinal stromal tumor: involvement in urologic patients and recent therapeutic advances. Urology. 2002;60(2):218-22.
58. Wingen CB, Pauwels PA, Debiec-Rychter M, van Gemert WG, Vos MC. Uterine gastrointestinal stromal tumour (GIST). Gynecologic oncology. 2005;97(3):970-2.

59. Takahashi RH, Matsubayashi J, Yokotsuka M, Tachibana M, Kusama H, Nagao T. An intrapelvic extraintestinal gastrointestinal stromal tumor of undetermined origin: diagnosis by prostate needle biopsy. Pathology, research and practice. 2012;208(12):736-40.

FIGURE/VIDEO CAPTIONS *figures should NOT be embedded in this document*

Fig. 1
Ultrasonographic image from dog 1 showing large round heterogeneous mass (between callipers) with multiple hypoechoic cavitations (asterisk shows the largest one) which was identified in the right cranial abdomen dorsal to the duodenum.

Fig. 2
Ultrasonographic image from dog 1 showing multiple different sized, hypoechoic nodules within the pancreatic parenchyma (between callipers) discovered five months after initial presentation. Note hepatic parenchyma in the near field (margin delineated with arrows).

Fig. 3
Abdominal gastrointestinal stromal tumour from dog 2 [HE]. The mass comprises a partially necrotic proliferation of moderately pleomorphic spindle cells arranged in interlacing streams.

Fig. 4
Immunohistochemistry of abdominal gastrointestinal stromal tumour from dog 2. Spindle cells label moderately strongly with antibody against CD117.

OWNER’S PERSPECTIVE *Optional*

Copyright Statement


I, *Joanna Lodzinska*, The Corresponding Author, has the right to assign on behalf of all authors and does assign on behalf of all authors, a full assignment of all intellectual property rights for all content within the submitted case report (other than as agreed with the BMJ Publishing Group Ltd and the British Veterinary Association) (“BMJ” and “BVA”) in any media known now or created in the future, and permits this case report (if accepted) to be published on Veterinary Record Case Reports and to be fully exploited within the remit of the assignment as set out in the assignment which has been read <http://journals.bmj.com/site/misc/vetreccrcopyright.pdf>

Date: 29/11/2018

PLEASE SAVE YOUR TEMPLATE WITH THE FOLLOWING FORMAT:


Corresponding author's last name and date of submission, eg,

Smith_June_2017.doc




Ultrasonographic image from dog 1 showing large round heterogeneous mass (between callipers) with multiple hypoechoic cavitations (asterisk shows the largest one) which was identified in the right cranial abdomen dorsal to the duodenum.

102x79mm (300 x 300 DPI)




Ultrasonographic image from dog 1 showing multiple different sized, hypoechoic nodules within the pancreatic parenchyma (between callipers) discovered five months after initial presentation. Note hepatic parenchyma in the near field (margin delineated with arrows).

115x77mm (300 x 300 DPI)



Abdominal gastrointestinal stromal tumour from dog 2 [HE]. The mass comprises a partially necrotic proliferation of moderately pleomorphic spindle cells arranged in interlacing streams.

59x44mm (600 x 600 DPI)



Immunohistochemistry of abdominal gastrointestinal stromal tumour from dog 2. Spindle cells label moderately strongly with antibody against CD117.

59x44mm (600 x 600 DPI)